

Effervescent Antacid & Pain Relief



National Brand Equivalent
Compare to Alka Seltzer®

Active Ingredients

- Aspirin (NSAID)
 - pain reliever
- Citric Acid
 - antacid
- Sodium Bicarbonate
 - antacid

Fast Relief of
Heartburn with Headache
Acid Indigestion
Upset Stomach
Body Aches & Pains

Standard Packouts 36 Count

- 2-count foil pouches (2 tabs per dose)
- 12 pack case,
 - 135 cases per pallet (ti/hi = 27/5)
- 72 pack case,
 - 20 cases per pallet (ti/hi = 5/4)

Minimum annual order quantity must be met in order to qualify for private label. We offer our house brand, Top Value, to those customers whose annual order quantities will not reach this threshold.

Drug Facts	
Active ingredients (in each effervescent tablet)	Purpose
Aspirin 325 mg (NSAID)*	Pain reliever
Citric acid 1000 mg	Antacid
Sodium bicarbonate (heat-treated) 1916 mg	Antacid
*Nonsteroidal anti-inflammatory drug	
Uses for the temporary relief of	
<ul style="list-style-type: none"> ■ heartburn, acid indigestion and sour stomach when accompanied by headache or body aches and pains ■ upset stomach with headache from overindulgence in food or drink ■ pain alone (headaches or body aches and pains) 	
Warnings	
<p>Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p> <p>Allergy alert: Aspirin may cause a severe allergic reaction which may include:</p> <ul style="list-style-type: none"> ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock <p>Stomach bleeding warning: This product contains a non-steroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed</p> <p>Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ if you are allergic to aspirin ■ if overly full from food or drink. Ask a doctor if severe stomach pain occurs after taking this product. ▶</p>	

Drug Facts (continued)													
<p>Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have a sodium-restricted diet</p> <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> ■ presently taking a prescription drug. Antacids may interact with certain prescription drugs. ■ taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis <p>When using this product do not take more than 8 tablets (adults) or 4 tablets (60 years and over) in a 24-hour period or use the maximum dosage for more than 10 days</p> <p>Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding</p> <ul style="list-style-type: none"> ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ an allergic reaction occurs. Seek medical help right away. ■ pain lasts for more than 10 days or gets worse ■ new symptoms occur ■ redness or swelling is present ■ ringing in the ears or a loss of hearing occurs <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>													
Directions													
<ul style="list-style-type: none"> ■ do not exceed recommended dosage ■ to avoid serious injury, completely dissolve tablets in 4 oz. of water before taking <table border="0"> <tr> <td>adults and children</td> <td>2 tablets every 4 hours</td> <td>do not exceed 8 tablets in 24 hours</td> </tr> <tr> <td>12 years and over</td> <td></td> <td></td> </tr> <tr> <td>(up to 60 years of age)</td> <td></td> <td></td> </tr> <tr> <td>60 years and over</td> <td>2 tablets every 4 hours</td> <td>do not exceed 4 tablets in 24 hours</td> </tr> </table>		adults and children	2 tablets every 4 hours	do not exceed 8 tablets in 24 hours	12 years and over			(up to 60 years of age)			60 years and over	2 tablets every 4 hours	do not exceed 4 tablets in 24 hours
adults and children	2 tablets every 4 hours	do not exceed 8 tablets in 24 hours											
12 years and over													
(up to 60 years of age)													
60 years and over	2 tablets every 4 hours	do not exceed 4 tablets in 24 hours											
Other information													
<ul style="list-style-type: none"> ■ each tablet contains: sodium 524 mg ■ store at room temperature (59°-86°F) 													
Inactive ingredients													
<ul style="list-style-type: none"> ■ none 													
<p>☎ Questions or comments? (telephone number)</p>													

Manufactured in the USA
421 110114

 **TOWER LABORATORIES LTD.**
YOUR PREMIER EFFERVESCENT SOURCE

8 Industrial Park Road • P.O. Box 306 • Centerbrook, CT 06409
8060 Whitbeck Road • Montague, MI 49437
www.towerlabs.com • Tel: (860) 767-2127 • Fax: (860) 767-2129