

Effervescent Antacid & Pain Relief



Extra Strength

Fast Relief of

Heartburn with Headache
Acid Indigestion
Upset Stomach
Body Aches & Pains

National Brand Equivalent
Compare to Alka Seltzer®

Active Ingredients

- Aspirin (NSAID)
 - pain reliever
- Anhydrous Citric Acid
 - antacid
- Sodium Bicarbonate
 - antacid

**Standard Packouts
36 Count**

- 2-count foil pouches (2 tabs per dose)
- 12 pack case,
 - 135 cases per pallet (ti/hi = 27/5)
- 72 pack case,
 - 20 cases per pallet (ti/hi = 5/4)

Minimum annual order quantity must be met in order to qualify for private label. We offer our house brand, Top Value, to those customers whose annual order quantities will not reach this threshold.

Drug Facts

Active ingredients (in each effervescent tablet)	Purpose
Anhydrous citric acid 1000 mg	Antacid
Aspirin 500 mg (NSAID)*	Pain reliever
Sodium bicarbonate (heat-treated) 1985 mg	Antacid

*Nonsteroidal anti-inflammatory drug

Uses for the temporary relief of

- heartburn, acid indigestion and sour stomach when accompanied by headache or body aches and pains
- upset stomach with headache from overindulgence in food or drink
- headache, body aches, and pain alone

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- if you are allergic to aspirin

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- presently taking a prescription drug. Antacids may interact with certain prescription drugs.
- taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis

When using this product do not take more than 8 tablets (adults) or 4 tablets (60 years and over) in a 24-hour period or use the maximum dosage for more than 2 weeks, except under the advice and supervision of a physician.

Drug Facts (continued)

Stop use and ask a doctor if you experience any of the following signs of stomach bleeding

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better

an allergic reaction occurs. Seek medical help right away.

- pain lasts for more than 10 days or gets worse
- new symptoms occur
- redness or swelling is present
- ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not exceed recommended dosage
- to avoid serious injury, completely dissolve tablets in 4 oz. of water before taking
- adults and children 12 years and over (up to 60 years of age) 2 tablets every 4 hours. Do not exceed 8 tablets in 24 hours.
- 60 years and over 2 tablets every 4 hours. Do not exceed 4 tablets in 24 hours.
- children under 12 years: consult a doctor

Other information

- each tablet contains: sodium 575 mg
- store at room temperature (59°-86°F)

Inactive ingredients:
flavor

(Questions or comments? (telephone number))

Manufactured in the USA
510 10/15/19



8 Industrial Park Road • P.O. Box 306 • Centerbrook, CT 06409
8060 Whitbeck Road • Montague, MI 49437
www.towerlabs.com • Tel: (860) 767-2127 • Fax: (860) 767-2129