Effervescent Pain Relief

National Brand Equivalent

Compare to Alka Seltzer®

Active Ingredients

Buffered aspirin (NSAID) - pain reliever & fever reducer

Standard Packout 20 Count

Minimum annual order quantity must be met in order to qualify for private label. We offer our house brand, Bromo Seltzer®, to those customers whose annual order quantities will not reach this threshold.

Fast Relief of

Headache Backache **Body Aches** Muscle Pain



Drug Facts

Active ingredient

Purpose

(in each effervescent tablet)

Buffered aspirin 325 mg (NSAID)*....Pain reliever/fever reduce *Nonsteroidal anti-inflammatory drug

Uses

- for the temporary relief of minor aches and pains due to
 - headache muscle pain backa
 toothache menstrual pain colds
 minor arthritis pain
- temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

■ hives ■ facial swelling ■ asthma (wheezing) ■ shoc

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn

you have high blood pressure, heart disease, liver

- **Drug Facts** (continued) sodium-restricted diet
 - cirrhosis, or kidney disease you are taking a diuretic you have asthma you have a

Ask a doctor or pharmacist before use if you are taking a prescription drug for

■ diabetes ■ gout ■ arthritis When using this product do not exceed recommended dosage

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding

- feel faint vomit blood have bloody or black stools
 have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away.
- pain lasts for more than 10 days or gets worse fever gets worse or lasts more than 3 days

- new symptoms occurredness or swelling is present
- ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not exceed recommended dosage
- to avoid serious injury, completely dissolve tablets in 4 oz. of water before taking

adults and children 12 years and over 2 tablets every 4 hours, or as directed by a doctor do not exceed 8 tablets in 24 hours 2 tablets every 4 hours, or as directed by a doctor do not exceed 4 tablets in 24 hours 60 years and over

children under 12 years consult a doctor

Other information

- each tablet contains: sodium 524 mg
 store at room temperature (59°-86°F). Avoid excessive heat

Inactive ingredients

anhydrous citric acid, sodium bicarbonate

(Questions or comments? (telephone number)

Manufactured in the USA with domestic and imported ingredients. Final label art to be determined at time of sale.

