

# Effervescent Antacid & Pain Relief

## Active Ingredients

Anhydrous Citric Acid  
- antacid  
Aspirin (NSAID)  
- pain reliever & fever reducer  
Sodium Bicarbonate  
- antacid

## Standard Packout 20 Count

Minimum annual order quantity must be met in order to qualify for private label. We offer our house brand, Bromo Seltzer®, to those customers whose annual order quantities will not reach this threshold.

**Fast Relief of**  
Acid Indigestion  
Upset Stomach  
Heartburn with Headache  
Body Aches & Pains



### Drug Facts

#### Active ingredients (in each effervescent tablet) Purpose

Anhydrous citric acid 1000 mg ..... Antacid  
Aspirin 325 mg (NSAID)\* ..... Pain reliever  
Sodium bicarbonate (heat-treated) 1916 mg ..... Antacid  
\*Nonsteroidal anti-inflammatory drug

#### Uses

for the temporary relief of  
■ heartburn, acid indigestion and sour stomach when accompanied by headache or body aches and pains  
■ upset stomach with headache from overindulgence in food or drink  
■ headache, body aches, and pain alone

#### Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

■ hives ■ facial swelling ■ asthma (wheezing) ■ shock  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Do not use** ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients.

**Ask a doctor before use if**  
■ stomach bleeding warning applies to you

### Drug Facts (continued)

■ you have a history of stomach problems, such as heartburn  
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease  
■ you are taking a diuretic ■ you have asthma ■ you have a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are** ■ presently taking a prescription drug. Antacids may interact with certain prescription drugs. ■ taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis

**When using this product** do not take more than 8 tablets (adults) or 4 tablets (60 years and over) in a 24-hour period or use the maximum dosage for more than 2 weeks, except under the advice and supervision of a physician.

**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding  
■ feel faint ■ vomit blood ■ have bloody or black stools  
■ have stomach pain that does not get better

■ an allergic reaction occurs. Seek medical help right away.  
■ pain lasts for more than 10 days or gets worse  
■ new symptoms occur  
■ redness or swelling is present  
■ ringing in the ears or a loss of hearing occurs

**If pregnant or breast-feeding,** ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

#### Directions

■ do not exceed recommended dosage  
■ to avoid serious injury, completely dissolve tablets in 4 oz. of water before taking

adults and children 12 years and over (up to 60 years of age)	2 tablets every 4 hours do not exceed 8 tablets in 24 hours
60 years and over	2 tablets every 4 hours do not exceed 4 tablets in 24 hours
Children under 12 years	consult a doctor

#### Other information

■ each tablet contains: sodium 524 mg  
■ store at room temperature (59°-86°F)

#### Inactive ingredients

■ none

**Questions or comments?** (telephone number)

Manufactured in the USA with domestic and imported ingredients.  
Final label art to be determined at time of sale.



**TOWER LABORATORIES LTD.**  
YOUR PREMIER EFFERVESCENT SOURCE

8 Industrial Park Road • P.O. Box 306 • Centerbrook, CT 06409  
www.towerlabs.com • Tel: (860) 767-2127 • Fax: (860) 767-2129