

Effervescent Antacid & Pain Relief

Extra Strength

Fast Relief of
Heartburn with Headache
Acid Indigestion
Upset Stomach
Body Aches & Pains



Active Ingredients

- Aspirin (NSAID)
 - pain reliever
- Anhydrous Citric Acid
 - antacid
- Sodium Bicarbonate
 - antacid

Standard Packout 24 Count

Minimum annual order quantity must be met in order to qualify for private label. We offer our house brand, Bromo Seltzer®, to those customers whose annual order quantities will not reach this threshold.

Drug Facts

Active ingredients (in each effervescent tablet)	Purpose
Anhydrous citric acid 1000 mg	Antacid
Aspirin 500 mg (NSAID)*	Pain reliever
Sodium bicarbonate (heat-treated) 1985 mg	Antacid

*Nonsteroidal anti-inflammatory drug

Uses for the temporary relief of

- heartburn, acid indigestion and sour stomach when accompanied by headache or body aches and pains
- upset stomach with headache from overindulgence in food or drink
- headache, body aches, and pain alone

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have a sodium-restricted diet
- you have asthma

Ask a doctor or pharmacist before use if you are

- presently taking a prescription drug. Antacids may interact with certain prescription drugs.
- taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis

When using this product do not take more than 8 tablets (adults) or 4 tablets (60 years and over) in a 24-hour period or use the maximum dosage for more than 2 weeks, except under the advice and supervision of a physician.

Drug Facts (continued)

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away.
- pain lasts for more than 10 days or gets worse
- new symptoms occur
- redness or swelling is present
- ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not exceed recommended dosage
- to avoid serious injury, completely dissolve tablets in 4 oz. of water before taking
- adults and children 12 years and over (up to 60 years of age) 2 tablets every 4 hours. Do not exceed 8 tablets in 24 hours.
- 60 years and over 2 tablets every 4 hours. Do not exceed 4 tablets in 24 hours.
- children under 12 years: consult a doctor

Other Information

- each tablet contains: sodium 575 mg
- store at room temperature (59°-86°F)

Inactive ingredients:
flavor

Questions or comments? (telephone number)

Manufactured in the USA with domestic and imported ingredients.
Final label art to be determined at time of sale.

 **TOWER LABORATORIES LTD.**
YOUR PREMIER EFFERVESCENT SOURCE

8 Industrial Park Road • P.O. Box 306 • Centerbrook, CT 06409
www.towerlabs.com • Tel: (860) 767-2127 • Fax: (860) 767-2129